


Relative Maturity:22

Positioning For:



MANAGEMENT COMMENTS

- Well suited for higher pH environments with known history of iron chlorosis deficiency.
- Solid option for early plant situations where SDS is not a concern.
- Canopy type suited to 30 inch rows or narrower.

SUITABILITY RATINGS

KEY ENVIRONMENTS

Field Prone to Lodging Highly Suitable

SUITABILITY

SCN-Prone Environments Highly Suitable
 SDS Prone Environments Poor Suitability
 High Residue Highly Suitable

SOILS

Early Planting/Cool Soils (w/SDS) Poor Suitability
 Drought-Prone Soils Poor Suitability
 Poorly Drained Soils Highly Suitable
 High PH Soils Highly Suitable

CHARACTERISTIC SCORES

Harvest Standability	8
Field Emergence	7
Phytoph. Field Tol.	4
Iron Def. Chlorosis	6
Canopy Width	5
Plant Height for Maturity	4
Relative Maturity	22
Technology Segment	RR2Y
Phytoph. Resist. Gene	1k
% Protein at 13% Moisture	32.7

DISEASE & PEST PROTECTION TRAITS

White Mold	4
Sudden Death Syndrome	3
SCN Race 1	4
Brown Stem Rot	8**
Aphid Antibiosis	A
Phytoph. Resist. Gene	
Iron Def. Chlorosis	

TRAIT SCORE RATINGS: 9 = Excellent; 1 = Poor. Canopy Width: 9 = Extremely Bushy; 1 = Very Narrow. Plant Height: 9 = Tall; 1 = Short. Blank = Insufficient Data. ** Ratings denoted with a double asterisk (**) reflect preliminary data subject to change when additional data becomes available.